arXiv:1203.6288v1 [math.NT] 28 Mar 2012

A CLOSED FORM EXPRESSION FOR THE DRINFELD MODULAR POLYNOMIAL $\Phi_T(X,Y)$

ALP BASSA AND PETER BEELEN

Abstract. In this paper we give a closed form expression for the Drinfeld modular polynomial $\Phi_T(X,Y) \in \mathbb{F}_q(T)[X,Y]$ for arbitrary q and prove a conjecture of Schweizer. A new identity involving the Catalan numbers plays a central role.

1. Introduction

Let $\mathbb{F}_q$ denote the finite field with q elements. For a polynomial $P(T) \in \mathbb{F}_q[T]$, the Drinfeld modular polynomial $\Phi_P(X,Y) \in \mathbb{F}_q(T)[X,Y]$ defines a model for the Drinfeld modular curve $X_0(P)$. Although the Drinfeld modular polynomials play an analogously fundamental role as the classical modular polynomials, very little is known about their explicit form. For levels P of low degree it is possible to compute $\Phi_P(X,Y)$ explicitly for relatively small values of q only. However, even in the case $P = T$ no explicit expression for $\Phi_T(X,Y)$ is known for general q. The main contribution of this paper is to find such an expression for general q by using a new identity involving Catalan numbers. Schweizer [6] studied $\Phi_T(X,Y)$, found an efficient algorithm to compute it for particular values of q and gave two conjectures concerning its structural properties. One of these conjectures was proven in [2], the second follows from a closed form expression for the Drinfeld modular polynomial $\Phi_T(X,Y)$ given below.

If ϕ is a Drinfeld module of rank two with j-invariant j and ϕ' is a P-isogenous Drinfeld module with j-invariant j', then $\Phi_P(j,j') = 0$. Moreover, the function field of $X_0(P)$ is given by $\mathbb{F}_q(T)(j,j')$. Since ϕ' is also P-isogenous to ϕ via the dual isogeny, the Drinfeld modular polynomial is symmetric. In case $P(T)$ is an irreducible polynomial, the extension degrees $[\mathbb{F}_q(T)(j,j') : \mathbb{F}_q(T)(j)]$ and $[\mathbb{F}_q(T)(j,j') : \mathbb{F}_q(T)(j')]$ are both equal to $q^{\deg P} + 1$.

If $\deg P = 1$, without loss of generality we may assume that $P = T$. The genus of the Drinfeld modular curve $X_0(P)$ is zero in this case. Schweizer [6] found an explicit relation between j, j' and a uniformizing parameter of the function field of $X_0(T)$. More precisely, he showed that the function field of $X_0(T)$ equals $\mathbb{F}_q(T)(z)$, where

$$j = \frac{(z+T)^{q+1}}{z} \text{ and } j' = \frac{(z+T^q)^{q+1}}{z^q}. \tag{1}$$

This implies that $\Phi_T(X,Y)$ is the unique monic, (symmetric) polynomial of degree $q+1$ in X (and Y), such that

$$\Phi_T\left(\frac{(z+T)^{q+1}}{z}, \frac{(z+T^q)^{q+1}}{z^q}\right) = 0. \tag{2}$$

2000 *Mathematics Subject Classification.* Primary: 11F32, 11F52, 05A10, 11B65; Secondary: 11G09, 14G35, 11F03.

Key words and phrases. Drinfeld Modular polynomial, Catalan Numbers, Drinfeld Modular Curves.

Peter Beelen was partially supported by DNRF (Denmark) and NSFC (China), grant No.11061130539.

As mentioned above, in [6] an algorithm to compute $\Phi_T(X,Y)$ was given. Also a start was made to describe $\Phi_T(X,Y)$ explicitly. More precisely, writing

$$\Phi_T(X,Y)=\sum_{m=0}^{q+1}P_m(Y)X^m$$

and defining $j_0=-T(T^{q-1}-1)^{q+1}$ it was shown that

- $P_{q+1}(Y)=1$
- $P_q(Y)=-(Y^q+T(Y-T^q)^{q-1}-T^{q^2}+T^q-T)$
- $P_1(Y)=-T^{q^2-2q+1}Y(Y-j_0)^{q-1}+(Y+T^{q^2}-T^q)(T^{q^3-q^2}-1)$
- $P_0(Y)=(Y-j_0)^{q+1}$

Moreover two conjectures were formulated concerning the structure of the polynomial $\Phi_T(X,Y)$. The first conjecture concerned a closed form expression for $\Phi_T(X,Y) \bmod T-1$, which was proven to hold in [2]. The second conjecture was the following:

Conjecture 1. *For all integers m satisfying $2\le m\le q-1$ we have*

$$\frac{P_m(Y)}{P_{q+1-m}(Y)}=\left(\frac{T^q(Y-j_0)}{Y-T^q}\right)^{q+1-2m}.$$

In this article we will state and prove a closed form expression for $\Phi_T(X,Y)$ (see Theorem 9). Conjecture 1 will be a direct consequence of the closed form expression (see Corollary 10). In the closed form the so-called Catalan numbers occur. These numbers are defined as

$$C_n:=\frac{1}{n+1}\binom{2n}{n}=\binom{2n}{n}-\binom{2n}{n-1} \tag{3}$$

and come up in a variety of combinatorial problems. In the next section we will show an identity involving Catalan numbers which is only valid in characteristic $p>0$. This identity will play an important role in Section 3, where a proof of the closed formula of $\Phi_T(X,Y)$ is given.

2. An identity involving the Catalan numbers

It will be useful to extend the definition of binomial coefficients $\binom{n}{k}$ to integers n and k:

$$\binom{n}{k}=\begin{cases}\dfrac{n(n-1)\cdots(n-k+1)}{k!} & \text{if } k\ge 0\\[2ex] 0 & \text{otherwise.}\end{cases} \tag{4}$$

The following lemma is a consequences of identities (5.21) and (5.43) in [5] and will be used later.

Lemma 1. *Let $r,m,n,$ and ℓ be integers and assume that $\ell\ge 0$. Then*

$$\binom{r}{m}\binom{m}{n}=\binom{r}{n}\binom{r-n}{m-n} \tag{5}$$

and

$$\sum_{i=0}^{\ell}(-1)^i\binom{r-i}{\ell}\binom{\ell}{i}=1. \tag{6}$$

With this lemma, we can prove the following:

Proposition 2. *Let $e \geq 2$ be a natural number. Then the following equality of polynomials in $\mathbb{Z}[t]$ holds:*

$$t^{e+1} - (-1)^e = \sum_{n=0}^{\lfloor \frac{e+1}{2} \rfloor} \left(\binom{e-n}{n-1} + \binom{e-n+1}{n} \right) (t-1)^{e+1-2n} t^n. \tag{7}$$

Proof. Using Newton's binomium to expand $(t-1)^{e+1-2n}$, we can rewrite equation (7):

$$\begin{aligned} t^{e+1} - (-1)^e &= \sum_{n=0}^{\lfloor \frac{e+1}{2} \rfloor} \sum_{m=0}^{e+1-2n} (-1)^{e+1-m} \left(\binom{e-n}{n-1} + \binom{e-n+1}{n} \right) \binom{e+1-2n}{m} t^{n+m} \\ &= \sum_{n \geq 0} \sum_{m \geq 0} (-1)^{e+1+m} \left(\binom{e-n}{n-1} + \binom{e-n+1}{n} \right) \binom{e+1-2n}{m} t^{n+m} \\ &= \sum_{n \geq 0} \sum_{i \geq n} (-1)^{e+1+i-n} \left(\binom{e-n}{n-1} + \binom{e-n+1}{n} \right) \binom{e+1-2n}{i-n} t^{i} \\ &= \sum_{i \geq 0} (-1)^{e+1+i} \sum_{n=0}^{i} (-1)^n \left(\binom{e-n}{n-1} + \binom{e-n+1}{n} \right) \binom{e+1-2n}{i-n} t^{i} \\ &= \sum_{i=0}^{e+1} (-1)^{e+1+i} \sum_{n=0}^{i} (-1)^n \left(\binom{e-n}{n-1} + \binom{e-n+1}{n} \right) \binom{e+1-2n}{i-n} t^{i}. \end{aligned}$$

Finally we see by moving the terms t^{e+1} and $(-1)^e$, that the proposition is equivalent to the identity

$$0 = \sum_{i=1}^{e} (-1)^{e+1+i} \sum_{n=0}^{i} (-1)^n \left(\binom{e-n}{n-1} + \binom{e-n+1}{n} \right) \binom{e+1-2n}{i-n} t^{i}. \tag{8}$$

At this point, note that by equation (5) we have:

$$\binom{e-n}{n-1}\binom{e+1-2n}{i-n} = \binom{e-n}{i-1}\binom{i-1}{n-1} \quad \text{and} \quad \binom{e-n+1}{n}\binom{e+1-2n}{i-n} = \binom{e-n+1}{i}\binom{i}{n}.$$

Further, by equation (6), we have for any $i \geq 1$:

$$\sum_{n=0}^{i} (-1)^n \binom{e-n}{i-1}\binom{i-1}{n-1} = -1 \quad \text{and} \quad \sum_{n=0}^{i} (-1)^n \binom{e-n+1}{i}\binom{i}{n} = 1.$$

Combining the above, we see that equation (8) and therefore the proposition, holds. □

The following lemma was shown in [2]:

Lemma 3. *Let $q = p^e$ be a power of a prime p and let n and k be integers satisfying $n \geq 0$ and $n - k < q$,. Then*

$$\binom{n}{k} \equiv (-1)^{n-k} \binom{q-1-k}{n-k} \pmod{p}.$$

This lemma has the following consequence:

Lemma 4. *Let i be an integer satisfying $0 \leq i < q-1$ and denote by C_i the i-th Catalan number. Then*

$$C_i \equiv (-1)^i \left(\binom{q-1-i}{i} + \binom{q-i}{i+1} \right) \pmod{p}. \tag{9}$$

Proof. Using equation (3) and Lemma 3, we find that

$$C_i = \binom{2i}{i} - \binom{2i}{i-1} \equiv (-1)^i \binom{q-1-i}{i} - (-1)^{i+1} \binom{q-i}{i+1} \pmod{p}.$$

The result now follows. □

We are now ready to state and prove an identity involving Catalan numbers that will turn out to be useful later.

Theorem 5. *Let q be a power of a prime p. Then*

$$\sum_{i=0}^{\lfloor \frac{q-1}{2} \rfloor} C_i \cdot (t(1-t))^i \equiv t^{q-1} + (1-t)^{q-1} \pmod{p}. \tag{10}$$

Proof. From equation (7) we see that (in characteristic zero)

$$\left(\frac{t}{t-1} \right)^{e+1} + \left(\frac{-1}{t-1} \right)^{e+1} = \sum_{i=0}^{\lfloor \frac{e+1}{2} \rfloor} \left(\binom{e-i}{i-1} + \binom{e-i+1}{i} \right) \left(\frac{t}{(t-1)^2} \right)^i.$$

Replacing t by $t/(t-1)$ we find

$$t^{e+1} + (1-t)^{e+1} = \sum_{i=0}^{\lfloor \frac{e+1}{2} \rfloor} (-1)^i \left(\binom{e-i}{i-1} + \binom{e-i+1}{i} \right) (t(1-t))^i.$$

Choosing $e = q$ and using Lemma 4, we find

$$t^{q+1} + (1-t)^{q+1} \equiv 1 - \sum_{i=1}^{\lfloor \frac{q+1}{2} \rfloor} C_{i-1} \cdot (t(1-t))^i \pmod{p}.$$

The theorem follows directly from this. □

Remark 6. *Note that Lemma 4 implies that $C_i \equiv 0 \pmod{p}$ for all i such that $(q-1)/2 < i < q-1$. This phenomenon has been observed in for example* [1].

Remark 7. *Note that t and $1-t$ are exactly the roots of the polynomial $s^2 - s + t(1-t)$. A reformulation of equation (10) is therefore:*

$$\sum_{i=0}^{\lfloor \frac{q-1}{2} \rfloor} C_i \cdot t^i \equiv s_1^{q-1} + s_2^{q-1} \pmod{p}, \tag{11}$$

where s_1 and s_2 are the two roots of the polynomial $s^2 - s + t$.

Remark 8. *The generating function of the Catalan numbers $c(x) := \sum_{i=0}^{\infty} C_i x^i$ can be expressed as*

$$c(x) = \frac{2}{1 + \sqrt{1-4x}}.$$

This implies that:

$$c\left(t(1-t)\right) = \frac{1}{1-t} = 1 + t + t^2 + \cdots.$$

Since $c(x)$ has integer coefficients, we can reduce this relation modulo a prime number p and deduce that $c\left(t(1-t)\right) \equiv 1/(1-t) \pmod{p}$. Equation (10) is a refined version of this, since it can be rewritten as

$$\sum_{i=0}^{\lfloor \frac{q-1}{2} \rfloor} C_i \cdot \left(t(1-t)\right)^i \equiv 1 + t + \cdots + t^{q-2} + 2t^{q-1} \pmod{p}.$$

Letting q tend to infinity, but holding p fixed, one recovers that $c\left(t(1-t)\right) \equiv 1/(1-t) \pmod{p}$.

3. A CLOSED FORM EXPRESSION FOR $\Phi_T(X,Y)$

Now we come to our main result:

Theorem 9. *Let q be a power of a prime p and define $j_0 = -T(T^{q-1}-1)^{q+1}$. Then*

$$\begin{aligned} \Phi_T(X,Y) &= (X+Y-j_0)^{q+1} - XY^q - X^qY + (XY)^q(T^{1-q}-1) + XY(T^{q-1}-1)^{q^2} \\ &\quad -T^{1-q}XY \sum_{i=0}^{\lfloor \frac{q-1}{2} \rfloor} C_i \cdot (XY - T^q(X+Y-j_0))^{q-1-2i}(XYT^{q^2+1})^i. \end{aligned} \tag{12}$$

Note that the term $(XY)^qT^{1-q}$ is canceled by a term in the summation corresponding to $i=0$. In fact, after expanding all terms in equation (12), one obtains a polynomial in X and Y with coefficients in $\mathbb{F}_q[T]$. Note that X^{q+1} and Y^{q+1} will have coefficient 1.

Proof. We will use the characterisation of $\Phi_T(X,Y)$ given in equation (2) and show that substituting

$$X = \frac{(z+T)^{q+1}}{z} \quad \text{and} \quad Y = \frac{(z+T^q)^{q+1}}{z^q} \tag{13}$$

on the left-hand side in equation (12), one obtains zero.

Equation (11) implies that

$$\sum_{i=0}^{\lfloor \frac{q-1}{2} \rfloor} C_i \cdot \left(\frac{XYT^{q^2+1}}{(XY - T^q(X+Y-j_0))^2} \right)^i = s_1^{q-1} + s_2^{q-1}, \tag{14}$$

with s_1 and s_2 equal to the two solutions to the equation

$$s^2 - s + \frac{XYT^{q^2+1}}{(XY - T^q(X+Y-j_0))^2} = 0. \tag{15}$$

Using equation (13), one can show after some computations that

$$XY - T^q(X+Y-j_0) = \frac{(T^{q^2+1}(z+T)^{q-1} + z^{q+1}(z+T^q)^{q-1})(z+T)(z+T^q)}{z^{q+1}} \tag{16}$$

and

$$\frac{XYT^{q^2+1}}{(XY - T^q(X+Y-j_0))^2} = \frac{T^{q^2+1}z^{q+1}(z+T)^{q-1}(z+T^q)^{q-1}}{(T^{q^2+1}(z+T)^{q-1} + z^{q+1}(z+T^q)^{q-1})^2}. \tag{17}$$

Equation (17) implies that the solutions s_1 and s_2 to equation (15) are given by

$$s_1 = \frac{T^{q^2+1}(z+T)^{q-1}}{T^{q^2+1}(z+T)^{q-1} + z^{q+1}(z+T^q)^{q-1}} \text{ and } s_2 = \frac{z^{q+1}(z+T^q)^{q-1}}{T^{q^2+1}(z+T)^{q-1} + z^{q+1}(z+T^q)^{q-1}}. \tag{18}$$

Using equations (13), (14), (16) and (18), we find that

$$\begin{aligned}
& T^{1-q}XY\sum_{i=0}^{\lfloor\frac{q-1}{2}\rfloor} C_i\cdot(XY-T^q(X+Y-j_0))^{q-1-2i}(XYT^{q^2+1})^i \\
= \quad & \\
& T^{1-q}XY(XY-T^q(X+Y-j_0))^{q-1}\sum_{i=0}^{\lfloor\frac{q-1}{2}\rfloor} C_i\cdot\left(\frac{XYT^{q^2+1}}{(XY-T^q(X+Y-j_0))^2}\right)^i \\
= \quad & \\
& T^{1-q}\frac{(z+T)^{2q}(z+T^q)^{2q}}{z^{q^2+q}}\left((T^{q^2+1}(z+T)^{q-1})^{q-1}+(z^{q+1}(z+T^q)^{q-1})^{q-1}\right) \\
= \quad & \\
& \frac{T^{q^3-q^2}(z+T)^{q^2+1}(z+T^q)^{2q}}{z^{q^2+q}}+\frac{(z+T)^{2q}(z+T^q)^{q^2+1}}{T^{q-1}z^{q+1}}.
\end{aligned}$$

However, a straightforward computation shows that, still with X and Y as in equation (13),

$$\begin{aligned}
& (X+Y-j_0)^{q+1}-XY^q-X^qY+(XY)^q(T^{1-q}-1)+XY(T^{q-1}-1)^{q^2} \\
= \quad & \\
& \frac{T^{q^3-q^2}(z+T)^{q^2+1}(z+T^q)^{2q}}{z^{q^2+q}}+\frac{(z+T)^{2q}(z+T^q)^{q^2+1}}{T^{q-1}z^{q+1}}.
\end{aligned}$$

This finishes the proof. □

We will now prove Conjecture 1 as a corollary to Theorem 9.

Corollary 10. *Define for* $0\le m\le q+1$ *polynomials* $P_m(Y)\in\mathbb{F}_q[T,Y]$ *by the identity*

$$\Phi_T(X,Y)=\sum_{m=0}^{q+1}P_m(Y)X^m.$$

Then for all m *such that* $2\le m\le q-1$ *we have*

$$\frac{P_m(Y)}{P_{q+1-m}(Y)}=\left(\frac{T^q(Y-j_0)}{Y-T^q}\right)^{q+1-2m}.$$

Proof. Note that the expression $(X+Y-j_0)^{q+1}-XY^q-X^qY+(XY)^q(T^{1-q}-1)+XY(T^{q-1}-1)^{q^2}$ only contributes to $P_m(Y)$ if $m\in\{0,1,q,q+1\}$. Now let m be an arbitrary integer between 2 and $q-1$. Since $XY-T^q(X+Y-j_0)=X(Y-T^q)-T^q(Y-j_0)$, we see from Theorem 9 that the coefficient of X^m in $\Phi_T(X,Y)$ is given by

$$\begin{aligned}
P_m(Y) &= T^{1-q}Y\sum_{i=0}^{\lfloor\frac{q-1}{2}\rfloor} C_i\cdot\binom{q-1-2i}{m-1-i}(Y-T^q)^{m-1-i}(-T^q(Y-j_0))^{q-m-i}(T^{q^2+1}Y)^i \\
&= T^{1-q}Y\left(\frac{Y-T^q}{-T^q(Y-j_0)}\right)^m\cdot\sum_{i=0}^{\lfloor\frac{q-1}{2}\rfloor} C_i\cdot\binom{q-1-2i}{m-1-i}\frac{(-T^q(Y-j_0))^{q-i}(T^{q^2+1}Y)^i}{(Y-T^q)^{1+i}}.
\end{aligned}$$

Since

$$\binom{q-1-2i}{m-1-i}=\binom{q-1-2i}{(q+1-m)-1-i},$$

we see that

$$\frac{P_m(Y)}{P_{q+1-m}(Y)}=\frac{((Y-T^q)/(-T^q(Y-j_0)))^m}{((Y-T^q)/(-T^q(Y-j_0)))^{q+1-m}}.$$

The corollary now follows directly. ☐

Remark 11. *Using Lemma 4 and Theorem 9, it is possible to give a closed form expression for* $\Phi_T(X,Y)$ *not involving the Catalan numbers. However, the resulting expression is not more compact nor easier to prove, as far as the authors know.*

Remark 12. *Setting* $t = x^{q-1}/(1+x^{q-1}) \in \mathbb{F}_q(x)$ *in equation (10) one obtains after some manipulations an algebraic dependency between* $u = x^q + x \in \mathbb{F}_q(x)$ *and* $v = x^{q+1}/(x^q+x) \in \mathbb{F}_q(x)$*, namely*

$$u^q \equiv u^{q-1}v^q + v + v \sum_{i=0}^{\lfloor \frac{q-1}{2} \rfloor} C_i u^{q-1-i} v^i \pmod{p}.$$

The functions u *and* v *occur in a natural way when defining a certain asymptotically optimal tower* [4]. *This tower was in* [3] *shown to have a Drinfeld modular interpretation. This again shows that equation (10) plays a role in the theory of Drinfeld modular curves.*

REFERENCES

[1] R. Alter and K.K. Kubota, Prime and prime power divisibility of Catalan numbers, Journal of Combinatorial Theory Series A 15, pp. 243-256 (1973).

[2] A. Bassa and P. Beelen, A proof of a conjecture by Schweizer on the Drinfeld modular polynomial $\Phi_T(X,Y)$, Journal of Number Theory 131, pp. 1276–1285 (2011).

[3] N.D. Elkies, Explicit towers of Drinfeld modular curves, Progress in Mathematics 202, pp. 189–198 (2001).

[4] A. Garcia and H. Stichtenoth, On the asymptotic behaviour of some towers of function fields over finite fields, Journal of Number Theory 61, pp. 248–273 (1996).

[5] R.L. Graham, D.E. Knuth and O. Patashnik, Concrete Mathematics, Addison-Wesley Publishing Company, 1989.

[6] A. Schweizer, On the Drinfeld Modular Polynomial $\Phi_T(X,Y)$, Journal of Number Theory 52, pp. 53–68 (1995).